



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

---

Forename(s)

---

Candidate signature

---

I declare this is my own work.

# A-level CHEMISTRY

## Paper 1 Inorganic and Physical Chemistry

Monday 10 June 2024

Morning

Time allowed: 2 hours

### Materials

For this paper you must have:

- the Periodic Table/Data Booklet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

### Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 105.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
<b>TOTAL</b>	



J U N 2 4 7 4 0 5 1 0 1

IB/M/Jun24/G4006/E9

**7405/1**

Answer **all** questions in the spaces provided.

0 1


This question is about atomic structure.

0 1 . 1

In 1897 JJ Thomson discovered the electron. He suggested that atoms were positively charged spheres with electrons embedded within them.

**Figure 1** represents an atom using Thomson's model.

**Figure 1**



Suggest the identity of this atom.

Give **two** differences between the modern model of an atom and the Thomson model of an atom.

**[3 marks]**

Identity \_\_\_\_\_

Difference 1 \_\_\_\_\_

\_\_\_\_\_

Difference 2 \_\_\_\_\_

\_\_\_\_\_



**0 1 . 2** Tellurium has a relative atomic mass of 127.6  
Iodine has a relative atomic mass of 126.9

Define relative atomic mass.

Suggest **one** property of tellurium that justifies its position before iodine in the modern Periodic Table.

**[3 marks]**

Definition \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Justification \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**0 1 . 3** A sample of tellurium is analysed in a time of flight (TOF) mass spectrometer using electron impact ionisation.

Give an equation, including state symbols, for this ionisation.

**[1 mark]**

\_\_\_\_\_

**Question 1 continues on the next page**

**Turn over ►**



0 1 . 4

In the TOF mass spectrometer an ion of an isotope of tellurium, with mass number  $y$ , travels along a 1.25 m flight tube with a kinetic energy of  $1.88 \times 10^{-12}$  J

The ion takes  $3.00 \times 10^{-7}$  s to reach the detector.

$$KE = \frac{1}{2} mv^2$$

$KE$  = kinetic energy / J

$m$  = mass / kg

$v$  = speed /  $\text{m s}^{-1}$

Calculate the mass, in g, of 1 mole of these tellurium ions.

Use your answer to suggest the mass number  $y$  of the tellurium isotope.

The Avogadro constant,  $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

[5 marks]

Mass \_\_\_\_\_ g

Mass number  $y$  \_\_\_\_\_



0 1 . 5

Tellurium has several other isotopes.  
Two of these isotopes are  $^{126}\text{Te}$  and  $^{124}\text{Te}$   
A different sample of tellurium is analysed using a TOF mass spectrometer.

Which statement about kinetic energy ( $KE$ ) is correct?

**[1 mark]**

Tick (✓) **one** box.

The  $KE$  of  $^{126}\text{Te}^+$  is greater than the  $KE$  of  $^{124}\text{Te}^+$

The  $KE$  of  $^{126}\text{Te}^+$  is the same as the  $KE$  of  $^{124}\text{Te}^+$

The  $KE$  of  $^{126}\text{Te}^+$  is less than the  $KE$  of  $^{124}\text{Te}^+$

13

**Turn over for the next question**

**Turn over ►**

**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**0 2**

This question is about an experiment to determine the solubility of strontium hydroxide in water at 20 °C

Strontium hydroxide is slightly soluble in water. Strontium hydroxide solution reacts in a similar way to calcium hydroxide solution.

- Some solid strontium hydroxide is added to approximately 1 dm<sup>3</sup> of distilled water in a stoppered flask.
- The mixture is kept at 20 °C. Every day, the mixture is checked. If no solid is present in the flask, more solid strontium hydroxide is added.
- On the day when no more solid needs to be added, the flask is opened and the mixture is filtered into another flask and stoppered.
- A 25.0 cm<sup>3</sup> sample of the filtrate is transferred to a conical flask with a pipette and a few drops of indicator added.
- This sample is titrated with 0.100 mol dm<sup>-3</sup> hydrochloric acid.
- The titration is repeated several times with further samples of the filtrate. The results are shown in **Table 1** on page 8.

**0 2 . 1**

Suggest why the solution is kept until no more solid needs to be added.

**[1 mark]**

---

---

**0 2 . 2**

Suggest why it is important to remove the undissolved strontium hydroxide before the titration.

**[1 mark]**

---

---

**0 2 . 3**

After the filtration, the solution is stored in a stoppered flask.

Suggest a reason for stoppering the flask.

**[1 mark]**

---

---

**Question 2 continues on the next page**

**Turn over ►**


**0 2 . 4** The diagrams in **Figure 2** show the part of a pipette with the graduation line.

Which diagram identifies the pipette that is correctly filled?

[1 mark]

Tick (✓) **one** box.

**Figure 2**



**0 2 . 5** Solubility can be quoted as 'g of solute per 100 cm<sup>3</sup> of solution'.

**Table 1** shows the results of the titrations between strontium hydroxide and hydrochloric acid. These can be used to determine the solubility of strontium hydroxide.

**Table 1**

Titration	Rough	1	2	3
Final burette reading / cm <sup>3</sup>	34.40	38.00	41.05	37.00
Initial burette reading / cm <sup>3</sup>	0.00	5.55	8.05	4.60
Titre / cm <sup>3</sup>	34.40	32.45	33.00	32.40

Give the equation for the reaction between strontium hydroxide and hydrochloric acid.

Use the results in **Table 1** to calculate the mean titre.

Use the mean titre to calculate the solubility of strontium hydroxide, in g per 100 cm<sup>3</sup> of solution, at 20 °C

[6 marks]



Equation

---

Mean titre \_\_\_\_\_ cm<sup>3</sup>

Solubility of strontium hydroxide \_\_\_\_\_ g per 100 cm<sup>3</sup> solution

_____
10

**Turn over for the next question**

**Turn over ►**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**0 3**

This question is about aqueous ions of the metal iron.

When an aqueous  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  ion reacts with ethanedioate ions, an iron(III) complex ion **X** is formed.

The only ligands in **X** are ethanedioate ions.

**0 3 . 1**

Draw the structure of **X**.

Include the charge.

**[2 marks]****0 3 . 2**

The formation of **X** is an example of the chelate effect.

Explain the meaning of the chelate effect.

**[2 marks]**

---

---

---

---

---

**Question 3 continues on the next page**

**Turn over ►**





**0 3 . 4** A student adds dilute ammonia solution to a solution containing  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  ions.

Give the formula of the precipitate that forms.

[1 mark]

---

**0 3 . 5** The student adds sodium carbonate solution to a solution containing  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  ions.

State **one** observation the student would make.

Give an equation for the reaction.

[2 marks]

Observation \_\_\_\_\_

Equation \_\_\_\_\_

---

**0 3 . 6** A solution containing  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  ions changes to a yellow-brown colour after several hours in contact with air.

The student adds sodium carbonate to the yellow-brown solution.

Give an equation for the reaction with sodium carbonate.

[1 mark]

---


14



**0 4**

This question is about some gas mixtures at equilibrium.

This reaction can be used to make hydrogen.

**0 4 . 1**

A mixture of 2.00 mol of  $\text{H}_2\text{O}(\text{g})$  and 2.00 mol of  $\text{CO}(\text{g})$  is allowed to reach equilibrium at a constant temperature in a  $20 \text{ dm}^3$  container.

At equilibrium, there are 0.92 mol of  $\text{H}_2(\text{g})$ .

Calculate the mole fraction of  $\text{H}_2(\text{g})$  in the equilibrium mixture.

**[2 marks]**

Mole fraction of  $\text{H}_2(\text{g})$  \_\_\_\_\_

**0 4 . 2**

State why the equilibrium constant ( $K_p$ ) for this reaction has no units.

**[1 mark]**

---

---

---

**Question 4 continues on the next page**

**Turn over ►**

0 4 . 3

The temperature of the equilibrium mixture formed in Question 04.1 is increased.

How does the amount of  $\text{H}_2(\text{g})$  change when the new position of equilibrium is reached?

**[1 mark]**


Tick (✓) **one** box.

The amount decreases.

The amount does not change.

The amount increases.

Ethanol can be made from ethene and steam.



**Table 2** shows the mole fractions of each of the gases in an equilibrium mixture at 6000 kPa

**Table 2**

Gas	Mole fraction
Ethene	0.645
Steam	0.323
Ethanol	0.0321



**0 4 . 4** Give an expression for  $K_p$  for this reaction.

Calculate the value of  $K_p$  at 6000 kPa

State the units.

**[4 marks]**

$K_p$

Units \_\_\_\_\_

**0 4 . 5** State the effect, if any, of an increase in volume of the container on the value of  $K_p$  for this reaction at a constant temperature.

**[1 mark]**

---

---

9

**Turn over for the next question**

**Turn over ►**



**0 5**

This question is about chlorine.

**0 5 . 1**

Give an equation to show how chlorine forms an acidic solution in water.

**[1 mark]**

---

**0 5 . 2**

Give an equation for the reaction between chlorine and cold, dilute aqueous sodium hydroxide.

**[1 mark]**

---

**0 5 . 3**In acidic conditions,  $\text{ClO}_3^-$  ions oxidise  $\text{Cl}^-$  ions to form  $\text{Cl}_2$ Deduce a half-equation for the oxidation of  $\text{Cl}^-$  to  $\text{Cl}_2$ Deduce a half-equation for the reduction of  $\text{ClO}_3^-$  to  $\text{Cl}_2$ 

Deduce the overall equation for this reaction.

**[3 marks]**Half-equation for the oxidation of  $\text{Cl}^-$  to  $\text{Cl}_2$ 

---

Half-equation for the reduction of  $\text{ClO}_3^-$  to  $\text{Cl}_2$ 

---

Overall equation

---



**0 5 . 4** Give the equation for the reaction of solid sodium chloride with concentrated sulfuric acid.

State the role of the chloride ions in this reaction.

**[2 marks]**

Equation

\_\_\_\_\_

Role

\_\_\_\_\_

**0 5 . 5** Draw the shape of the  $\text{Cl}_3^-$  ion.  
Include any lone pairs of electrons that influence the shape.

**[1 mark]**

**0 5 . 6** Chlorine forms an ion with the Group 3 element thallium (Tl).

State and explain the bond angle in  $\text{TlCl}_2^+$

**[2 marks]**

Bond angle

\_\_\_\_\_

Explanation

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

10

Turn over ►



0 6

This question is about vanadium ions.

**Table 3** shows some standard electrode potential values.

**Table 3**

	$E^\ominus / \text{V}$
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00
$\text{VO}^{2+}(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{V}^{3+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.34
$\text{V}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{V}^{2+}(\text{aq})$	-0.26
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$\text{V}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{V}(\text{s})$	-1.20
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.38

0 6 . 1

Use the data in **Table 3** to explain why Zn reduces an aqueous solution of  $\text{VO}_2^+$  ions to  $\text{V}^{2+}$  ions, but does not reduce it any further.

[2 marks]

---



---



---



---

0 6 . 2

Identify the species in **Table 3** that can reduce an aqueous solution of  $\text{VO}_2^+$  to V

[1 mark]

---



**0 6 . 3** Two half-cells  $\text{Fe}^{2+}(\text{aq}) / \text{Fe}(\text{s})$  and  $\text{VO}^{2+}(\text{aq}) / \text{V}^{3+}(\text{aq})$  are connected.

Calculate the EMF of this cell.

Give the conventional representation for this cell.

Give a half-equation for the reaction that occurs at the negative electrode.

**[3 marks]**

EMF \_\_\_\_\_

Cell representation

---

Half-equation

---

**Question 6 continues on the next page**

**Turn over ►**




0 6 . 4

0.151 g of impure  $\text{NH}_4\text{VO}_3$  is added to dilute sulfuric acid to form a solution containing aqueous  $\text{VO}_2^+$  ions.

All the  $\text{VO}_3^-$  ions are converted to  $\text{VO}_2^+$  ions.

These  $\text{VO}_2^+$  ions are reduced to aqueous  $\text{V}^{2+}$  ions by reaction with an excess of zinc.




The excess of zinc is removed by filtration and washed.

The filtrate, containing the  $\text{V}^{2+}$  ions, is titrated with a  $0.0200 \text{ mol dm}^{-3}$  solution of acidified  $\text{KMnO}_4$

$29.43 \text{ cm}^3$  of  $\text{KMnO}_4$  solution are needed to oxidise all the  $\text{V}^{2+}$  ions to  $\text{VO}_2^+$  ions.

The ionic equation for the reaction of  $\text{MnO}_4^-$  ions with  $\text{V}^{2+}$  ions is



Calculate the percentage purity of the  $\text{NH}_4\text{VO}_3$

Give your answer to 3 significant figures.

[4 marks]

Percentage purity \_\_\_\_\_

10



**0 7**At 40 °C the ionic product of water,  $K_w = 2.92 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ **0 7 . 1**Give the expression for  $K_w$ Calculate the pH of pure water at 40 °C  
Give your answer to 2 decimal places.**[3 marks]** $K_w$ 

pH \_\_\_\_\_

**0 7 . 2**35.0 cm<sup>3</sup> of 0.150 mol dm<sup>-3</sup> aqueous sodium hydroxide are mixed with  
20.0 cm<sup>3</sup> of a 0.100 mol dm<sup>-3</sup> solution of hydrochloric acid.  
The temperature of the solution formed is 40 °CCalculate the pH of the solution formed.  
Give your answer to 2 decimal places.**[5 marks]**

pH \_\_\_\_\_

**8**

Turn over ►



0 8

This question is about enthalpy changes.

0 8 . 1

Theoretical values for enthalpies of lattice dissociation can be calculated using a perfect ionic model.

State the meaning of the term perfect ionic model.

[1 mark]

---




---

0 8 . 2

Enthalpies of lattice dissociation can also be obtained from Born–Haber cycles.

**Figure 3** shows an incomplete Born–Haber cycle for the formation of sodium oxide.

**Figure 3**



Complete **Figure 3** by writing formulas, including state symbols, of the appropriate species on each of the two blank lines.

[2 marks]



**0 8 . 3** Table 4 shows some enthalpy changes.

**Table 4**

Enthalpy change	$\Delta H / \text{kJ mol}^{-1}$
Enthalpy of atomisation of oxygen	+248
Enthalpy of atomisation of sodium	+109
Enthalpy of formation of sodium oxide	-416
First ionisation energy of sodium	+494
First electron affinity of oxygen	-142
Second electron affinity of oxygen	+844

Use the data in **Table 4** to calculate the enthalpy of lattice dissociation of sodium oxide.

**[2 marks]**

Enthalpy of lattice dissociation \_\_\_\_\_  $\text{kJ mol}^{-1}$

**0 8 . 4** Explain why the second electron affinity of oxygen has a positive value.

**[1 mark]**

---

---

---

**Question 8 continues on the next page**

**Turn over ►**



0 8 . 5

Explain why the enthalpy of lattice dissociation for sodium oxide is greater than the enthalpy of lattice dissociation for sodium chloride.

**[2 marks]**

---

---

---

---

0 8 . 6

Sodium chloride dissolves in water.

**Table 5** shows some more enthalpy changes.

**Table 5**

Enthalpy change	$\Delta H / \text{kJ mol}^{-1}$
Enthalpy of hydration for $\text{Cl}^-$ ions	-364
Enthalpy of hydration for $\text{Na}^+$ ions	-406
Enthalpy of lattice dissociation for NaCl	+771

Use the data in **Table 5** to calculate the enthalpy of solution for sodium chloride.

**[2 marks]**

Enthalpy of solution \_\_\_\_\_  $\text{kJ mol}^{-1}$



0 8 . 7

Give a reason why data books do **not** contain a value for the enthalpy of solution of sodium oxide.


**[1 mark]**

---

---

0 8 . 8

Calculate the temperature, in °C, at which this reaction becomes feasible.

**[3 marks]**

Temperature \_\_\_\_\_ °C

14

**Turn over for the next question****Turn over ►**

**0 9**

This question is about metals and their compounds.

**0 9 . 1**

State why the atomic radius of calcium is greater than the atomic radius of magnesium.

**[1 mark]**

---

---

---

**0 9 . 2**

Magnesium reacts with steam.

Give an equation, including state symbols, for this reaction.

**[1 mark]**

---



0	9	.	3
---	---	---	---

Similar-sized pieces of barium and magnesium are added to separate 100 cm<sup>3</sup> samples of dilute sulfuric acid. In each case the sulfuric acid is in excess.

The barium reacts quickly at first. After a few minutes the reaction stops, even though there is still some unreacted barium in the flask.

The magnesium reacts more slowly than the barium, but the reaction continues until all the magnesium has reacted.

Explain why

- the barium initially reacts more quickly than the magnesium
- the barium reaction stops before all the barium has reacted.

**[3 marks]**

---

---

---

---

---

---

---


**Question 9 continues on the next page**

**Turn over ►**



0	9	.	4
---	---	---	---

A metal nitrate  $\text{X}(\text{NO}_3)_2$  completely decomposes when heated.



A 0.832 g sample of  $\text{X}(\text{NO}_3)_2$  decomposes on heating to produce a total of 348 cm<sup>3</sup> of gas at 298 K and 100 kPa

Deduce the identity of metal **X**.

The ideal gas constant,  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

**[6 marks]**

Identity of metal **X** \_\_\_\_\_



**0 9 . 5** Sodium reacts with aluminium and hydrogen to form solid  $\text{NaAlH}_4$

Give an equation for this reaction.

Suggest why  $\text{NaAlH}_4$  has a high melting point.

**[3 marks]**

Equation

---

Suggestion

---

---

---

**0 9 . 6** Give the equation for the reaction between  $\text{H}_3\text{PO}_4$  and an excess of  $\text{NaOH}$

**[1 mark]**

---

Lithium is an important metal used in cells to power mobile phones.

**0 9 . 7** In a lithium cell, a lithium cobalt oxide electrode and a lithium electrode are used.

Give the equation for the reaction that occurs at the positive electrode.

**[1 mark]**

---

**0 9 . 8** Commercial electrochemical cells can be rechargeable or non-rechargeable.

State why lithium cells can be recharged.

**[1 mark]**

---

---

**END OF QUESTIONS**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**









**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk).

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2024 AQA and its licensors. All rights reserved.



3 6



2 4 6 A 7 4 0 5 / 1

IB/M/Jun24/7405/1