



A-level
CHEMISTRY
7405/2

Paper 2 Organic and Physical Chemistry

Mark scheme

June 2024

Version: 1.1 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

No student should be disadvantaged on the basis of their gender identity and/or how they refer to the gender identity of others in their exam responses.

A consistent use of 'they/them' as a singular and pronouns beyond 'she/her' or 'he/him' will be credited in exam responses in line with existing mark scheme criteria.

Further copies of this mark scheme are available from aqa.org.uk

Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2024 AQA and its licensors. All rights reserved.

AS and A-Level Chemistry

Mark Scheme Instructions for Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the examiner make his or her judgement and help to delineate what is acceptable or not worthy of credit or, in discursive answers, to give an overview of the area in which a mark or marks may be awarded.

The extra information in the 'Comments' column is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

You should mark according to the contents of the mark scheme. If you are in any doubt about applying the mark scheme to a particular response, consult your Team Leader.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which might confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

The use of M1, M2, M3 etc in the right-hand column refers to the marking points in the order in which they appear in the mark scheme. So, M1 refers to the first marking point, M2 the second marking point etc.

2. Emboldening

- 2.1** In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **OR**. Different terms in the mark scheme are shown by a / ;eg allow smooth / free movement.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general 'List' principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (often prefaced by 'Ignore' in the mark scheme) are not penalised.

For example, in a question requiring 2 answers for 2 marks:

| Correct answers | Incorrect answers (i.e. incorrect rather than neutral) | Mark (2) | Comment |
|-----------------|--|----------|---|
| 1 | 0 | 1 | |
| 1 | 1 | 1 | They have not exceeded the maximum number of responses so there is no penalty. |
| 1 | 2 | 0 | They have exceeded the maximum number of responses so the extra incorrect response cancels the correct one. |
| 2 | 0 | 2 | |
| 2 | 1 | 1 | |
| 2 | 2 | 0 | |
| 3 | 0 | 2 | The maximum mark is 2 |
| 3 | 1 | 1 | The incorrect response cancels out one of the two correct responses that gained credit. |
| 3 | 2 | 0 | Two incorrect responses cancel out the two marks gained. |
| 3 | 3 | 0 | |

3.2 Marking procedure for calculations

Full marks should be awarded for a correct numerical answer, without any working shown, unless the question states 'Show your working' or 'justify your answer'. In this case, the mark scheme will clearly indicate what is required to gain full credit.

If an answer to a calculation is incorrect and working is shown, process mark(s) can usually be gained by correct substitution / working and this is shown in the 'Comments' column or by each stage of a longer calculation.

3.3 Errors carried forward, consequential marking and arithmetic errors

Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ECF or consequential in the marking scheme.

An arithmetic error should be penalised for one mark only unless otherwise amplified in the marking scheme. Arithmetic errors may arise from a slip in a calculation or from an incorrect transfer of a numerical value from data given in a question.

3.4 Equations

In questions requiring students to write equations, state symbols are generally ignored unless otherwise stated in the 'Comments' column.

Examiners should also credit correct equations using multiples and fractions unless otherwise stated in the 'Comments' column.

3.5 Oxidation states

In general, the sign for an oxidation state will be assumed to be positive unless specifically shown to be negative.

3.6 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.7 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term or if the question requires correct IUPAC nomenclature.

3.8 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.9 Ignore / Insufficient / Do not allow

Ignore or insufficient is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

Do **not** allow means that this is a wrong answer which, even if the correct answer is given, will still mean that the mark is not awarded.

3.10 Marking crossed out work

Crossed out work that **has not been** replaced should be marked as if it were not crossed out, if possible. Where crossed out work **has been** replaced, the replacement work and not the crossed out work should be marked.

3.11 Reagents

The command word "Identify", allows the student to choose to use **either** the name or the formula of a reagent in their answer. In some circumstances, the list principle may apply when both the name and the formula are used. Specific details will be given in mark schemes.

The guiding principle is that a reagent is a chemical which can be taken out of a bottle or container. Failure to identify complete reagents **will be penalised**, but follow-on marks (e.g. for a subsequent equation or observation) can be scored from an incorrect attempt (possibly an incomplete reagent) at the correct reagent. Specific details will be given in mark schemes.

For example, **no credit** would be given for

- the cyanide ion or CN^- when the reagent should be potassium cyanide or KCN;
- the hydroxide ion or OH^- when the reagent should be sodium hydroxide or NaOH;

- the $\text{Ag}(\text{NH}_3)_2^+$ ion when the reagent should be Tollens' reagent (or ammoniacal silver nitrate). In this example, no credit is given for the ion, but credit could be given for a correct observation following on from the use of the ion. Specific details will be given in mark schemes.

In the event that a student provides, for example, **both** KCN and cyanide ion, it would be usual to ignore the reference to the cyanide ion (because this is not contradictory) and credit the KCN. Specific details will be given in mark schemes.


3.12 Organic structures

Where students are asked to draw organic structures, unless a specific type is required in the question and stated in the mark scheme, these may be given as displayed, structural or skeletal formulas or a combination of all three as long as the result is unambiguous.

In general

- Displayed formulae must show all of the bonds and all of the atoms in the molecule, but need not show correct bond angles.
- Skeletal formulae must show carbon atoms by an angle or suitable intersection in the skeleton chain. Functional groups must be shown and it is essential that all atoms other than C atoms are shown in these (except H atoms in the functional groups of aldehydes, secondary amines and N-substituted amides which do not need to be shown).
- Structures must not be ambiguous, e.g. 1-bromopropane should be shown as $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ and not as the molecular formula $\text{C}_3\text{H}_7\text{Br}$ which could also represent the isomeric 2-bromopropane.
- Bonds should be drawn correctly between the relevant atoms. This principle applies in all cases where the attached functional group contains a carbon atom, e.g nitrile, carboxylic acid, aldehyde and acid chloride. The carbon-carbon bond should be clearly shown. Wrongly bonded atoms will be penalised **on every occasion**. (see the examples below)
- The same principle should also be applied to the structure of alcohols. For example, if students show the alcohol functional group as $\text{C} - \text{HO}$, they should be penalised **on every occasion**.
- Latitude should be given to the representation of $\text{C} - \text{C}$ bonds in alkyl groups, given that CH_3- is considered to be interchangeable with $\text{H}_3\text{C}-$ even though the latter would be preferred.
- Similar latitude should be given to the representation of amines where $\text{NH}_2 - \text{C}$ will be allowed, although $\text{H}_2\text{N} - \text{C}$ would be preferred.
- Poor presentation of vertical $\text{C} - \text{CH}_3$ bonds or vertical $\text{C} - \text{NH}_2$ bonds should **not** be penalised. For other functional groups, such as $-\text{OH}$ and $-\text{CN}$, the limit of tolerance is the half-way position between the vertical bond and the relevant atoms in the attached group.

By way of illustration, the following would apply.



- Representation of CH₂ by C–H₂ will be penalised
- Some examples are given here of **structures** for specific compounds that should **not** gain credit (but, exceptions may be made in the context of balancing equations)

CH₃COH for ethanal

CH₃CH₂HO for ethanol

OHCH₂CH₃ for ethanol

C₂H₆O for ethanol

CH₂CH₂ for ethene

CH₂.CH₂ for ethene

CH₂:CH₂ for ethene

- Each of the following **should gain credit** as alternatives to correct representations of the structures.

CH₂ = CH₂ for ethene, H₂C=CH₂

CH₃CHOHCH₃ for propan-2-ol, CH₃CH(OH)CH₃

- In most cases, the use of “sticks” to represent C – H bonds in a structure should **not** be penalised. The exceptions to this when “sticks” will be penalised include
 - structures in mechanisms where the C – H bond is essential (e.g. elimination reactions in halogenoalkanes and alcohols)
 - when a displayed formula is required
 - when a skeletal structure is required or has been drawn by the candidate

3.13 Organic names

As a general principle, non-IUPAC names or incorrect spelling or incomplete names should **not** gain credit. Some illustrations are given here.


Unnecessary but not wrong numbers will **not** be penalised such as the number ‘2’ in 2-methylpropane or the number ‘1’ in 2-chlorobutan-1-oic acid.

| | |
|-------------------------|--|
| but-2-ol | should be butan-2-ol |
| 2-hydroxybutane | should be butan-2-ol |
| butane-2-ol | should be butan-2-ol |
| 2-butanol | should be butan-2-ol |
| ethan-1,2-diol | should be ethane-1,2-diol |
| 2-methylpropan-2-ol | should be 2-methylpropan-2-ol |
| 2-methylbutan-3-ol | should be 3-methylbutan-2-ol |
| 3-methylpentan | should be 3-methylpentane |
| 3-methylpentane | should be 3-methylpentane |
| 3-methylpentane | should be 3-methylpentane |
| propanitrile | should be propanenitrile |
| aminethane | should be ethylamine (although aminoethane can gain credit) |
| 2-methyl-3-bromobutane | should be 2-bromo-3-methylbutane |
| 3-bromo-2-methylbutane | should be 2-bromo-3-methylbutane |
| 3-methyl-2-bromobutane | should be 2-bromo-3-methylbutane |
| 2-methylbut-3-ene | should be 3-methylbut-1-ene |
| difluorodichloromethane | should be dichlorodifluoromethane |


3.14 Organic reaction mechanisms

Curly arrows should originate either from a lone pair of electrons or from a bond.

The following representations should not gain credit **and will be penalised each time** within a clip.



For example, the following would score zero marks



When the curly arrow is showing the formation of a bond to an atom, the arrow can go directly to the relevant atom, alongside the relevant atom or **more than half-way** towards the relevant atom.

In free-radical substitution

- the absence of a radical dot should be penalised **once only** within a clip.
- the use of half-headed arrows is not required, but the use of double-headed arrows or the incorrect use of half-headed arrows in free-radical mechanisms should be penalised **once only** within a clip

The correct use of skeletal formulae in mechanisms is acceptable, but where a C-H bond breaks, both the bond and the H must be drawn to gain credit.

3.15 Extended responses

For questions marked using a 'Levels of Response' mark scheme:

Level of response mark schemes are broken down into three levels, each of which has a descriptor. Each descriptor contains two statements. The first statement is the Chemistry content statement and the second statement is the communication statement.

Determining a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the Chemistry content descriptor for that level. The descriptor for the level indicates the qualities that might be seen in the student's answer for that level. If it meets the lowest level, then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer.

When assigning a level you should look at the overall quality of the answer and not look to pick holes in small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Once the level has been decided, the mark within the level is determined by the communication statement:

- If the answer completely matches the communication descriptor, award the higher mark within the level.
- If the answer does not completely match the communication descriptor, award the lower mark within the level.

The exemplar materials used during standardisation will help you to determine the appropriate level. There will be an exemplar in the standardising materials which will correspond with each level of the mark scheme and for each mark within each level. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the exemplar to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the exemplar.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.


Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other chemically valid points. Students may not have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme. The mark scheme will state how much chemical content is required for the highest level.

An answer which contains nothing of relevance to the question must be awarded no marks.

For other extended response answers:

Where a mark scheme includes linkage words (such as 'therefore', 'so', 'because' etc), these are optional. However, a student's marks for the question may be limited if they do not demonstrate the ability to construct and develop a sustained line of reasoning which is coherent, relevant, substantiated and logically structured. In particular answers in the form of bullet pointed lists may not be awarded full marks if there is no indication of logical flow between each point or if points are in an illogical order.

The mark schemes for some questions state that the maximum mark available for an extended response answer is limited if the answer is not coherent, relevant, substantiated and logically structured. During the standardisation process, the Lead Examiner will provide marked exemplar material to demonstrate answers which have not met these criteria. You should use these exemplars as a comparison when marking student answers.

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--|------------------------|
| 01.1 | <p>Curve eg</p>  <p>[Br₂] / mol dm⁻³</p> <p>Time / s</p> | <p>Curve with decreasing gradient</p> <p>A straight line with a negative gradient was also allowed. (This is only appropriate if the concentrations of the other reagent are in such large excess that they are effectively constant, but, since the concentrations of the other reagents are not specified, credit was given for taking this approach.)</p> | <p>1 (1 x AO1)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|------------------------|
| 01.2 | <p>M1 For $[\text{OH}^-] = 7.50 \times 10^{-2}$</p> <p>M2 For rate = 2.75×10^{-11}</p> <p>M3 $k = \frac{\text{rate}}{[\text{CH}_3\text{COCH}_3][\text{OH}^-]}$</p> <p>OR $k = \frac{2.75 \times 10^{-11}}{(1.5 \times 10^{-2}) \times (2.5 \times 10^{-2})}$</p> <p>M4 $k = 7.3(3) \times 10^{-8}$</p> <p>M5 units = $\text{mol}^{-1}\text{dm}^3\text{s}^{-1}$</p> | <p>M3 For rearranging rate equation</p> <p>Or</p> <p>For inserting correct numbers in rearranged equation</p> <p>If rearrangement upside down lose M3 but can score M4 for 1.36×10^7 as ECF M5 for $\text{mol dm}^{-3} \text{ s}$ as ECF</p> | <p>5 (5 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|------------------------|
| 01.3 | <p>M1 Arrow from C–H bond to C–C</p> <p>M2 Arrow from C=O bond to O</p> <p>M3 Arrow from lone pair on O to C–O bond</p> <p>M4 Arrow from Br–Br bond to Br</p> | <p>Dipoles must be correct if shown for M4</p> | <p>4 (4 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|----------------|
| 01.4 | Step 1 includes CH_3COCH_3 and OH^- <u>and</u> these are also in the rate equation Or Step 1 contains all the species in the rate equation | Br_2 not in step 1 <u>and</u> not in rate equation so it has to be step 1 | 1 (1 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--------------------------------|----------------|
| 02.1 | M1 mol P = $0.0145 + (2 \times 0.0115) = 0.0375$ M2 $[P] = \frac{M1}{0.025} = 1.50 \text{ mol dm}^{-3}$ | ECF from incorrect M1 | 2 (2 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|----------------|
| 02.2 | M1 $K_c = \frac{[R][S]^3}{[P]^2 [Q]}$ M2 $K_c = \frac{\left(\frac{0.0115}{0.045}\right)\left(\frac{0.0345}{0.045}\right)^3}{\left(\frac{0.0145}{0.045}\right)^2 \left(\frac{0.0275}{0.045}\right)}$ or $= \frac{(0.256)(0.767)^3}{(0.322)^2(0.611)}$ M3 = 1.81 to 1.82 M4 units mol dm^{-3} | M1 Must be square brackets in expression M2 Inserts values and divides by volume in dm^3 M3 Evaluates expression If no use of volume lose M2 but can score M3 for 0.0817 M4 Allow consequential to their expression | 4 (4 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|---------------------------|
| 02.3 | <p>M1 <u>equilibrium</u> shifts to side with most moles</p> <p>M2 to oppose decrease in concentration of all reactants and products / dilution of everything</p> <p>OR</p> <p>M1 K_c is expressed as a function of concentrations and concentration equals amount over volume.</p> <p>M2 If Volume increases the amount of R and S must increase in order to keep K_c constant.</p> | <p>Allow</p> <p>M2 oppose the decrease in concentration of S</p> <p>$K_c = \frac{RS^3}{P^2Q} \times \frac{1}{V}$ (where R,S etc are amounts)</p> <p>So, if V increases R and S must increase relative to P and Q to keep K_c constant</p> | <p>2</p> <p>(2 x AO3)</p> |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--------------------------------|----------------|
| 03.1 | M1 + 3C ₂ H ₆ M2 Zeolite / Aluminosilicate / Aluminium oxide | | 2 (2 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|----------|--------------------------------|----------------|
| 03.2 | Option B | | 1 (1 x AO3) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---------|--------------------------------|----------------|
| 03.3 | Alkenes | | 1 (1 x AO1) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|-------------|---|---|----------------|
| 03.4 | M1 Initial volume O ₂ = 0.21 × 1350 = 283.5 (cm ³) M2 Volume of O ₂ remaining = M1 – (6.5 × 20) = 153.5 cm ³ M3 Volume of CO ₂ formed = 20 × 4 = 80 cm ³ M4 Total volume of gas left = M2 + M3 + (0.79 × 1350) = 1300 cm ³ | Alternative route: M1 Vol Air decreases by 6.5 × 20 = 130 cm ³ M2 = 1220 cm ³ M3 Vol CO ₂ produced = 4 × 20 = 80 cm ³ M4 Total Vol Air + CO ₂ = 1220 + 80 = 1300 cm ³ | 4 (4 × AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|-------------|---|---|----------------|
| 03.5 | M1 Acid rain M2 CaO or CaCO ₃ | M1 Allow damages (limestone) buildings or statues / death of aquatic organisms / air pollution | 2 (2 × AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|---------------------------------------|
| 04.1 | <p>M1 $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3 + \text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COOCH}(\text{CH}_3)_2 + \text{H}_2\text{O}$</p> <p>M2 Methyl ethyl ethanoate.</p> |  <p>Allow ECF from incorrect 5 carbon ester</p> <p>Allow other valid names 1-methylethyl ethanoate Isopropyl ethanoate 2-propyl ethanoate Propan-2-yl ethanoate</p> | <p>2 (1 x AO1, 1 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|-------------|---|--|------------------------|
| 04.2 | <p>This question is marked using Levels of Response. Refer to the Mark Scheme Instructions for Examiners for guidance.</p> | <p>Indicative Chemistry Content</p> <p>Stage 1</p> <p>1a Measuring cylinder(s) for the propan-2-ol and ethanoic acid (size not required but if specified should be between 10 - 100 cm³) Allow 10cm³ / graduated pipette or burette</p> <p>1b (Dropping/teat) pipette for sulfuric acid (NOT graduated or other qualification for pipette)</p> <p>Stage 2 Diagram eg below to include</p> <p>2a Labelled <u>condenser</u> shown vertical and open at top and bottom i.e. in cross section</p> <p>2b Labelled <u>water in</u> at the bottom and <u>water out</u> at the top of the vertical condenser</p> <p>2c Labelled reaction <u>flask</u> recognisable as either pear shaped or round bottomed</p> | <p>6 (6 x AO3)</p> |
| | <p>Level 3 5–6 marks</p> <p>All stages are virtually complete (virtually complete means one from stage 1 and two from stages 2 and 3)</p> <p>Answer communicates the whole explanation, including equations, coherently and shows a logical progression through all three stages</p> | | |
| | <p>Level 2 3–4 marks</p> <p>All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies (covered means one from a stage)</p> <p>OR two stages virtually complete (virtually complete means one from stage 1 or two from stages 2 and 3)</p> <p>Answer is coherent and shows some progression through all three stages. Some steps in each stage may be incomplete</p> | | |
| | <p>Level 1 1–2 marks</p> <p>Two stages are covered (covered means one from a stage) but the explanation of each stage may be incomplete or may contain inaccuracies</p> <p>OR only one stage is virtually complete (virtually complete means one from stage 1 or two from stages 2 and 3)</p> <p>Answer shows some progression between two stages</p> | | |

| | | | | |
|--|--|--|--|--|
| | | | <p>Stage 3 Safety Needs precaution AND reason for each suggestion</p> <p>3a Use a fume cupboard / fume hood / well-ventilated lab space AND to avoid breathing in harmful / toxic / corrosive compounds</p> <p>3b Wear gloves AND as compounds are corrosive</p> <p>3c Add glass beads/chips (to the mixture before heating) / labelled as anti-bumping granules/chips AND to ensure smooth boiling / reduce size of bubbles</p> <p>3d Use an electric heater/water bath AND as compounds are flammable</p> | |
|--|--|--|--|--|

| | | | | |
|-------------------------------|---|---|--|--|
| <p>04.2 (cont)</p> | <p>0 mark</p> | <p>Insufficient correct chemistry to gain a mark.</p> | | |
| | <div style="display: flex; justify-content: space-around; align-items: center;">  <div style="margin-left: 20px;"> <p>2a Condenser shown vertical and open at the top and bottom (cross sectional)</p> <p>2b Water labelled in at the bottom AND out at the top</p> <p>2c Reaction flask recognisable as EITHER pear shaped OR round bottomed</p> </div> </div> | | | |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|----------------------------|
| 04.3 | M1 To neutralise / remove / react with (excess) acid M2 Remove stopper/bung OR tip the funnel upside down and open the tap M3 There will be a build up of pressure / gas / carbon dioxide OR M2 Allow add stopper M3 To prevent spillage | M3 must be linked to their precaution in M2 | 3 (1 x AO1, 2 x AO3) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--------------------------------|--------------------------------|----------------|
| 04.4 | Drying agent / To remove water | Not dehydrating agent | 1 (1 x AO1) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|---|----------------|
| 04.5 | Compare boiling point to a data book/known value | Boils at sharp boiling point / over a narrow temp range | 1 (1 x AO1) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|----------------|
| 05.1 | M1 Misty / white / steamy fumes M2 No visible change | M1 (immediate) White precipitate forms M2 White precipitate forms <u>slowly</u> | 2 (2 x AO3) |






| Question | Answers | Additional Comments/Guidelines | Mark | | | | | | | | | | |
|----------|---|--|--|---|--|---|----|--|--|--|---|--|----------------|
| 05.2 | <p>M1 Propanal AND (blue solution gives a brick) red precipitate</p> <p>If M1 incorrect, allow ECF for suitable tests on remaining liquids</p> <table border="1"> <tbody> <tr> <td>M2</td> <td>(Warm with) acidified potassium dichromate (VI)</td> <td>add Na</td> <td>warm with a named carboxylic acid with conc H₂SO₄</td> <td>(Warm with) acidified potassium manganate (VII)</td> </tr> <tr> <td>M3</td> <td>Propan-1-ol / alcohol AND (orange solution) goes green</td> <td>Propan-1-ol/ alcohol AND effervescence</td> <td>Propan-1-ol / alcohol AND fruity smell</td> <td>Propan-1-ol / alcohol AND (purple solution) goes colourless</td> </tr> </tbody> </table> | M2 | (Warm with) acidified potassium dichromate (VI) | add Na | warm with a named carboxylic acid with conc H ₂ SO ₄ | (Warm with) acidified potassium manganate (VII) | M3 | Propan-1-ol / alcohol AND (orange solution) goes green | Propan-1-ol/ alcohol AND effervescence | Propan-1-ol / alcohol AND fruity smell | Propan-1-ol / alcohol AND (purple solution) goes colourless | | 3 (3 x AO3) |
| M2 | (Warm with) acidified potassium dichromate (VI) | add Na | warm with a named carboxylic acid with conc H ₂ SO ₄ | (Warm with) acidified potassium manganate (VII) | | | | | | | | | |
| M3 | Propan-1-ol / alcohol AND (orange solution) goes green | Propan-1-ol/ alcohol AND effervescence | Propan-1-ol / alcohol AND fruity smell | Propan-1-ol / alcohol AND (purple solution) goes colourless | | | | | | | | | |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--------------------------------|----------------|
| 06.1 | $\text{C}_4\text{H}_9\text{COOH} + \text{NaHCO}_3 \rightarrow \text{C}_4\text{H}_9\text{COONa} + \text{CO}_2 + \text{H}_2\text{O}$ | | 1 (1 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|----------------|
| 06.2 |  | $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{COOH}$ | 1 (1 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|----------------|
| 06.3 | M1  M2 6:1:2:1 (Any order) | M1 $(\text{CH}_3)_2\text{CHCH}_2\text{COOH}$ M2 Allow ECF for a 5 carbon carboxylic acid | 2 (2 x AO2) |



| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|------------------------|
| 06.4 | <p>M1</p>  <p>Or</p> <p>M2 Adjacent C has no (non-equivalent) H attached (so no splitting/spin-spin coupling takes place)</p> | <p>M1 $(\text{CH}_3)_2\text{C}(\text{OH})\text{COCH}_3$ or $\text{CH}_3\text{OC}(\text{CH}_3)_2\text{CHO}$</p> | <p>2 (2 x AO2)</p> |


| Question | Answers | | | | Additional Comments/Guidelines | Mark |
|----------|---|------------------|--------------|---|--------------------------------|----------------------------|
| 06.5 |  | Scores M1 and M2 | Allow M1 for |  | | 6 (1 x AO1, 5 x AO3) |
| |  | Scores M3 and M4 | Allow M3 for | RCH ₃ | | |
| |  | Scores M5 | | This structure also scores M5  | | |
| | ¹³ C peaks | = 3 | M6 | Allow ECF from their M5 of C ₅ H ₁₀ O ₂ | | |

| Question | Answers | Additional Comments/Guidelines | Mark | | | | | | | | | |
|-----------------------|---|---|------|---|---|-----------------------|-----------------------|-------------------------|---|---|---|--|
| 07 | <p>M1 $\frac{2.62 \times 10^{-3}}{6.56 \times 10^{-4}} = 4$ or $\frac{6.56 \times 10^{-4}}{2.62 \times 10^{-3}} = 0.25$</p> <p>M2 Hence $4\text{CO}_2 + 4\text{H}_2\text{O}$</p> <p>M3 So 4C and 8H in L</p> <p>M4 Hence 2O so $\text{C}_4\text{H}_8\text{O}_2$</p> <p>$\text{C}_3\text{H}_4\text{O}_3$ scores 1 if no other mark scored</p> | <p>Alternative method</p> <p>M1 $n\text{H in L} = 5.24 \times 10^{-3}$ Hence mass H = 5.24×10^{-3} g</p> <p>M2 $n\text{C in L} = 2.62 \times 10^{-3}$ Hence mass C = $2.62 \times 10^{-3} \times 12$ $= 3.144 \times 10^{-2}$ g</p> <p>M3 Mass L = $6.56 \times 10^{-4} \times 88$ $= 0.057728$ g mass O = $0.057728 - (5.24 \times 10^{-3} + 3.144 \times 10^{-2})$ $= 0.021048$ g</p> <p>M4 $n\text{O} = 0.021048 / 16$ $= 1.3155 \times 10^{-3}$</p> <p>EF</p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;">C</td> <td style="text-align: center;">H</td> <td style="text-align: center;">O</td> </tr> <tr> <td style="text-align: center;">2.62×10^{-3}</td> <td style="text-align: center;">5.24×10^{-3}</td> <td style="text-align: center;">1.3155×10^{-3}</td> </tr> <tr> <td style="text-align: center;">2</td> <td style="text-align: center;">4</td> <td style="text-align: center;">1</td> </tr> </table> <p>MF = $(88/44) \times \text{C}_2\text{H}_4\text{O}$</p> <p>= $\text{C}_4\text{H}_8\text{O}_2$</p> <p>$\text{C}_3\text{H}_4\text{O}_3$ scores 1 if no other mark scored</p> | C | H | O | 2.62×10^{-3} | 5.24×10^{-3} | 1.3155×10^{-3} | 2 | 4 | 1 | <p style="text-align: center;">4 (4 x AO2)</p> |
| C | H | O | | | | | | | | | | |
| 2.62×10^{-3} | 5.24×10^{-3} | 1.3155×10^{-3} | | | | | | | | | | |
| 2 | 4 | 1 | | | | | | | | | | |



| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|-------------------------------------|--------------------------------|----------------|
| 08.1 | M1 Electrophilic Addition M2 HBr | | 2 (2 x AO1) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--------------------------------|----------------------------|
| 08.2 | M1 Butan-2-ol or correct structure eg $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$ M2 NaOH AND (warm) aqueous (Allow cold if stated) | | 2 (1 x AO1, 1 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|-------------|--|--------------------------------|-------------------------------------|
| <p>08.3</p> | <p>M1 HCN or KCN/H₂SO₄</p>  <p>M2 Arrow from lone pair on C to C of C=O</p> <p>M3 Arrow from C=O to O</p> <p>M4 Structure of intermediate including negative on O</p> <p>M5 Arrow from lone pair on O to H⁺</p> <p>OR</p>  <p>M2</p> <p>M3</p> <p>M4</p> <p>M5</p> | | <p>5 (1 x AO1, 4 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|------------------------|
| 08.4 | <p>M1</p>  <p>planar/ flat</p> <p>M2 Equal chance (50/50) attack from above/below OWTTE</p> <p>M3 Giving equal amounts of both optical isomers/enantiomers</p> | <p>Allow planar carbonyl group</p> <p>Either side</p> | <p>3 (3 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|---|----------------------------|
| 09.1 | M1 CH_3NH_2 Shown as displayed or abbreviated structural formula M2 N-methyl ethylamine or N-methyl ethanamine For M2 allow alkyl groups reversed | For M2 Allow N-methyl aminoethane or N-methyl N-ethylamine or Methyl ethylamine Or Methyl ethanamine | 2 (1 x AO1, 1 x AO2) |


| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--------------------------------|----------------|
| 09.2 | <p>M1 </p> <p>M2 </p> <p>Note: If answers are non-skeletal penalise once only</p> | | 2 (2 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--------------------------------|------------------------|
| 09.3 | <p>M1 For structure of 2 bromo propane</p> $ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{Br} \\ \\ \text{H} \end{array} \quad \text{M1} $ <p>M2 For TWO correct curly arrows</p>  <p>Step 1</p> $ \begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{Br} \\ \\ \text{H} \end{array} + \text{:CN}^- \longrightarrow (\text{CH}_3)_2\text{CHCN} + \text{Br}^- $ <p>Step 2</p> $ (\text{CH}_3)_2\text{CHCN} + 2\text{H}_2 \longrightarrow \text{Amine G} $ <p>M3 Amine G</p> <p>M3 Amine G has a <u>fully displayed</u> structure of Amine G</p> | | <p>3 (3 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--------------------------------|----------------|
| 09.4 | M1 The <u>lone pair on nitrogen</u> in P is more available or more able to accept protons/H ⁺ M2 more alkyl groups are electron releasing/donating or greater (positive) inductive effect (of the alkyl groups) | | 2 (2 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|------------------------------------|
| 10.1 | <p>M1 Benzene is more stable than cyclohexatriene</p> <p>M2 The enthalpy of hydrogenation of benzene is (152 kJ mol^{-1}) less / less exothermic</p> <p>M3 Due to the delocalisation of electrons in benzene</p> <p>M4 Both are planar / hexagonal</p> <p>M5 Benzene has equal C-C bond lengths or regular hexagon whereas Cyclohexa-1,3,5-triene has bonds of different/varied length or the hexagon is distorted/irregular</p> | <p>M4 and M5 could be shown in a clear diagram</p> | <p>5 (2 x AO1 3 x AO3)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--------------------------------|---------------------------|
| 10.2 | M1 Concentrated nitric acid AND concentrated sulfuric acid / conc. HNO_3 AND conc. H_2SO_4 M2 $\text{HNO}_3 + 2\text{H}_2\text{SO}_4 \rightarrow \text{NO}_2^+ + \text{H}_3\text{O}^+ + 2\text{HSO}_4^-$ OR $\text{HNO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{NO}_3^+ + \text{HSO}_4^-$ then $\text{H}_2\text{NO}_3^+ \rightarrow \text{NO}_2^+ + \text{H}_2\text{O}$ OR $\text{HNO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{O} + \text{NO}_2^+ + \text{HSO}_4^-$ | | 2 (1 x AO1 1 x AO2) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|----------------|
| 10.3 |  <p>M1</p> <p>M2 – structure</p> <p>M3</p> | M1 Positive must be on N and arrow from inside hexagon to N or + on N M2 Structure showing horseshoe and positive charge. <ul style="list-style-type: none"> Horseshoe centred on C1 but must not extend beyond C2 and C6 + in intermediate not too close to C1 (allow on or “above” a line from C2 to C6) M3 Arrow from C-H bond back into hexagon | 3 (3 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--|----------------|
| 10.4 | Sn/HCl Ignore references to NaOH used after Sn/HCl BUT penalise if NaOH used at the same time as Sn/HCl | Allow H ₂ with Pt/Ni Allow HCl with Fe | 1 (1 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|---|--|----------------|
| 10.5 | <u>Manufacture</u> of dyes/(cationic) surfactants/fabric softener | Allow to <u>make</u> hair/fabric conditioner | 1 (1 x AO1) |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--|--|
| 11.1 | <p>M1 $Q = (mc \Delta T = 60 \times 4.18 \times 52.1) = 13066.68 \text{ J}$</p> <p>M2 moles = $(\frac{1.31}{136}) = 0.00963 \text{ mol}$</p> <p>M3 $\frac{Q}{n} = \frac{13066.68}{0.00963} = 1356541 \text{ (J mol}^{-1}\text{)}$</p> <p>M4 $\Delta H = -1360 \text{ kJ mol}^{-1}$</p> | <p>M3 = M1 / M2</p> <p>M4 = - M3 /1000</p> <p>Allow range -1355 to -1362</p> | <p>4</p> <p>(1 x AO1, 3 x AO2)</p> |

| Question | Answers | Additional Comments/Guidelines | Mark |
|----------|--|--|---------------------------|
| 11.2 | <p>Any 5 from the following:</p> <p>M1 Value from calorimetry less exothermic / lower / smaller (than method 1 value)</p> <p>M2 (Calorimetry =) because of heat / energy loss</p> <p>M3 (Calorimetry =) incomplete combustion</p> <p>M4 (Calorimetry =) some liquid hydrocarbon could have evaporated</p> <p>M5 Mean bond enthalpies values use enthalpies taken across a range of compounds</p> <p>M6 Value from bond enthalpy data ignores energy changes in vaporisation of the fuel or condensing the water</p> | <p>M1 Allow both are less (than method 1)</p> <p>M2 Copper absorbs some heat energy</p> <p>M6 Allow value from mean bond enthalpies does not include changes of state. M6 Allow value from mean bond enthalpies use gaseous states</p> | <p>5</p> <p>(5 x AO3)</p> |